

Stranger Danger? An investigation into the influence of human-horse bond on stress and behaviour

Ijichi, Carrie; Griffin, Kym; Squibb, Keith; Favier, Becky

Published in:
Applied Animal Behaviour Science

Publication date:
2018

The re-use license for this item is:
CC BY-NC-ND

This document version is the:
Peer reviewed version

The final published version is available direct from the publisher website at:
[10.1016/j.applanim.2018.05.034](https://doi.org/10.1016/j.applanim.2018.05.034)

Find this output at Hartpury Pure

Citation for published version (APA):
Ijichi, C., Griffin, K., Squibb, K., & Favier, B. (2018). Stranger Danger? An investigation into the influence of human-horse bond on stress and behaviour. *Applied Animal Behaviour Science*, 206(September).
<https://doi.org/10.1016/j.applanim.2018.05.034>

Stranger Danger? An investigation into the influence of human-horse bond on stress and behaviour.

Carrie Ijichi ^{1*}, Kym Griffin², Keith Squibb ¹ & Rebecca Favier¹

¹ *University Centre Hartpury, Hartpury, Gloucester, UK, GL19 3BE*

² *Nottingham Trent University, College of Science and Technology, School of Animal Rural & Environmental Sciences, Burton Street, Nottingham, UK NG1 4BU*

*email: carrie.ijichi@hartpury.ac.uk

*Tel: 01452 702618

ABSTRACT

Human-animal bond is receiving increasing attention and is thought to confer benefits on well-being and performance in working animals. One important benefit of bonding is the “safe base” an attachment figure provides, which manifests in better coping and increased exploration during potential threat. However, there is limited research exploring the existence or benefits of human-horse bonds, though bonding is sought after by both pleasure and elite riders. The purpose of the current study was to determine whether the presence of horses’ owners confers a safe-base, improving horse behaviour and physiological stress responses during novel handling tests. Horses completed two different handling tests, one with their owner and the other with an unfamiliar experimental handler (n = 46). Test and handler order was randomised and handlers were double blind to the performance of the horse with the alternate handler. Time taken to complete the tests and proactive behaviour were measured as indicators of performance and compliance. Core temperature, discrepancy in eye temperature, heart rate and heart rate variability were recorded to assess stress responses. If horses experience a “safe base” effect in the vicinity of their owner, they would be expected to show lower stress responses and greater behavioural compliance, compared to being handled by a stranger. However, there was no difference in behaviour or any physiological stress response between the handlers. This indicates that a calm, competent, but unknown handler is equally effective to an owner during stressful procedures as neither equine performance nor affective state supported a safe-base effect. This supports previous research suggesting that the level of bond between human and horse is not the most salient factor in equine well-being or compliance during training and handling. These findings have implications for veterinary and clinical behaviour counselling, where novel human handlers must modify behaviour under potentially stressful circumstances.

KEY TERMS: infrared thermography; heart rate variability; bond; trust; horse; handling;

1. INTRODUCTION

Human-animal bond has received increasing interest in recent years (e.g. Payne et al. 2016; Payne et al. 2015). Attachment Theory is concerned with the development of bonds between infants and their caregivers in humans (Cassidy, 1999) and mammalian species (Newberry and Swanson, 2008). It is theorised that appropriate bonds aid in survival because vulnerable offspring keep close to their mothers in such species. Since domestic animals depend on human caregivers to a certain extent, some level of attachment-type bond may exist. A fully developed relationship bond is characterised by proximity seeking, secure base, safe haven and separation distress (Cassidy, 1999). Secure base refers to reduced stress under perceived threat and increased exploration in the presence of the attachment figure (Mikulincer and Shaver, 2003). It is therefore, a suitable construct of bonding to investigate objectively in human-animal bonds.

Bonding between animals and their human caregivers is highly desirable as it is purported to improve human well-being (Walsh, 2009) and is anecdotally reported to affect training outcomes in horses (e.g. Parelli 1993; Roberts 1997). Within competitive equestrianism, human-horse bonds are thought to be integral to the success of partnerships during challenging and highly pressurised situations (Fallis, 2013). However, due to this perceived importance, and the fact that many human carers feel strong bonds towards their animal companions, it may be that reciprocal bonds are incorrectly perceived. Species that are highly dependent upon their caregiver, such as dogs, may be presumed to have more opportunities to bond. Indeed, separation anxiety is a relatively commonly recognised phenomenon in dogs isolated from their owners (Riemer et al., 2016) and the safe base effect has been observed in dogs (INSERT MIKLOSI).

Horses do not live as inter-dependently with their carers, yet studies indicate that horses can discern the difference between familiar and unfamiliar humans and that this elicits different cognitive responses (Proops and McComb, 2012). Therefore, it is possible that such bonds do form in a species that does not live in such close proximity with their carers, though this has not yet been investigated to our knowledge.

Whilst familiarity is known to have positive influences on behaviour during handling in horses (Marsbøll and Christensen, 2015), the effect of more complex bonds that may result from longer term interactions has not been assessed. Therefore, the current study aims to determine whether horses respond differently to novelty, depending on whether they are with their owner or a stranger. To this end, horses completed two novel handling tests, one with their owner and the other with an unknown experimental handler. Time taken to complete the task and proactivity during refusal were measured as indicators of compliance and performance. Heart rate, heart rate variability, core temperature and the discrepancy between eye temperatures were measured as physiological indicators of stress and affective states. If an owner provides a safe base as the result of a human-horse bond (Cassidy, 1999), horses would be expected to take less time to complete the tasks, show less potentially dangerous proactive behaviour and have lower physiological indicators of stress, compared to when handled by an unfamiliar person.

2. METHOD

The current experiment was conducted within an indoor arena at Hartpury College Equestrian Centre, Gloucestershire (UK) in October 2016. Subjects were liveries at this facility which allowed testing to occur in a home arena, reducing the effects of environmental novelty (Wolff et al., 1997). Forty-six horses of mixed breeds and genders (26 geldings and 20 mares) took part. Age ranged from 3 – 20 years (mean = 9.33 ± 4.20). All subjects had completed at least

preliminary work under saddle. Subjects were housed and managed as per owner preferences on a large livery yard. In general, subjects were provided forage three times a day with hard-feed dependent on workload and nutritional requirements and constant access to fresh water. They were individually stabled with a minimum of 1 hour of exercise each day but with limited or no turn-out at the time of testing. The typical method of training was not known and will depend on owner preference, temperament and knowledge. Therefore, subjects are likely to have been trained differently regarding positive and negative reinforcement. Subjects were handled in their own headcollar, providing it did not include inbuilt pressure mechanisms.

2.1 Handlers

The familiar handler was the owner and daily care-giver of the subject. The unfamiliar handler was the same for all subjects (C.I.) and had not made contact with any subject prior to testing. This individual was a competent, experienced handler and had completed similar handling tests before (Ijichi et al., 2013). The experimental handler wore the same clothing for all tests, whilst owners were free to choose their own attire. This was to reduce the potential effect of clothing on how subjects perceived the unfamiliar handler (Hausberger et al., 2008). Both the owner and experimental handler wore gloves, a riding helmet and protective footwear.

2.2 Handling Tests

Tests required subjects to navigate novel objects in response to leadrope pressure, which is an aid used to indicate that the horse should step forward (McGreevy and McLean, 2007). Each test was sufficiently different to prevent habituation, which might alter behaviour between the first and the second test. Task A consisted of a 2.5m x 3m blue tarpaulin secured to the surface of the indoor holding arena by 20 individual tent pegs (Ijichi et al., 2013). To complete this test, the subject walked over the tarpaulin. Test B consisted of a frame that was 2.5m high and 1.6m wide, from which hung 2m long coloured plastic streamers (Squibb et al., 2018). To complete

99 this test, the subject walked through the frame, causing the streamers to touch the face and
100 body of the subject as they passed through.

101 Both objects were present within the test arena and faced the exit and conspecifics, because
102 differing directions could have affected the motivation to complete the test. A standard jump
103 pole was placed 2m in front of each test, which the subject walked over to mark the start of
104 the test. Handlers indicated that the horse should walk towards the obstacle using leadrope
105 pressure but no verbal or additional tactile cues were permitted. Horses had a maximum of 3
106 minutes to complete each handling test, as previous research indicates that horses that have
107 not completed the test within this time do not do so (Ijichi et al., 2013). Tests were recorded on
108 video for post-hoc analysis.

109 *2.3 Experimental Design*

110 Upon arrival at the testing area, horses were fitted with a Polar Equine V800 heart rate monitor
111 by K.G. (Polar Electro Oy, Kempele, Finland). The elasticated surcingle was attached to the
112 girth area, which had been moistened with water to aid conductivity. After confirming that HR
113 was being detected, subjects were given a minimum of 5 minutes to habituate to the monitor.
114 This was deemed sufficient as all subjects had previously worn girths and/or lunging rollers.
115 During habituation, subjects were outside of the indoor testing arena and could not see the
116 novel objects.

117 Test order and handler order was randomised and horse order was pseudo-randomised,
118 depending on the availability of subjects. Each handler was blind to the behaviour of the subject
119 with the alternate handler and owners were expressly forbidden from discussing the likely
120 behaviour of the subject. Double-blinding was possible as the test arena had solid doors and a
121 research assistant remained outside at all times to prevent the second handler from attempting
122 to see into the arena. Subjects entered the arena with the first handler and proceeded to a
123 designated area for eye temperature measurement. This was marked by two parallel jump poles

in the same position and direction within the enclosed area. This was to reduce the potentially confounding effects of direct sunlight and environmental factors on IRT readings (Church et al., 2014). The research assistant (K.S.) stood at a marked point approximately 1m and 90 degrees from each eye (Travain et al., 2015; Yarnell et al., 2013). Images were taken using a FLIR E4 thermal imaging camera (FLIR Systems, USA.). The handler then led the subject towards Test A or B as randomly allocated.

Upon successful completion of the task, or termination at 3 minutes, the subject was led back to the designated area for post-test eye temperature readings. Recordings were taken as per pre-test procedures. Horses that completed the task in less than 3 minutes were then held for the remainder of the available crossing time. This ensured the second handler could not deduce the subject's behaviour during the first task as all horses remained in the arena for a similar amount of time. Upon leaving the test arena, the subject had a minimum of 5 minutes to recover, before re-entering with the second handler. The procedure was then repeated verbatim.

2.4 Analysis

2.4.1 Behaviour

Crossing time began when the first fore-limb bore weight after the ground pole 2m in front of the obstacle. Crossing time ended when the last hind-limb bore weight on the tarpaulin for Test A (Ijichi et al., 2013), or when the tail of the subject had passed through the frame for Test B (Squibb et al., 2018). Horses that did not complete the test were recorded a Crossing Time of 180 seconds. Proactivity (outlined below) was calculated as per Ijichi et al. (2013). Refusal behaviour was defined as any behaviour which did not contribute to crossing the object. This included moving backwards, sideways, forwards but away from the object, rearing or remaining stationary. Refusal that lasted for 10 seconds or more was analysed to determine how proactive that refusal was (Tarpaulin: N = 13, Streamers: N = 36). Proactive refusal was defined as any refusal behaviour that involved movement thus excluding stationary refusal. Proactive refusal

was then recorded as the percent of total refusal time for any individual which showed refusal behaviour (which included remaining stationary). A higher value indicated a greater amount of proactive behaviour (Ijichi et al., 2013). Twelve subjects exhibited refusal behaviour for both tests, allowing a comparison between handlers.

2.4.2 Infrared Thermography

IRT was analysed using FLIR Tools software (ver. 5.9.16284.1001) post-hoc. This was to reduce any stress inducing effects of prolonged IRT recordings (Travain et al., 2015) required to record accurate readings from a small area. Eye temperature recordings were the maximum temperature within the palpebral fissure from the lateral commissure to the lacrimal caruncle (Yarnell et al., 2013). A mean of the left and right eyes was calculated for each subject, pre and post-test, for each test. In addition, the temperature of the left eye was subtracted from the right eye to indicate the discrepancy between both eyes, pre and post-test, for each test. A positive score indicates a hotter right eye, whilst a negative score indicates a hotter left eye. This may provide an indicator of ipsilateral hemispheric dominance (Lush and Ijichi, 2018) and lateralised processing of stimuli (De Boyer Des Roches et al., 2008).

2.4.3 Heart Rate

Heart rate readings were taken from the point of the first IRT reading to the second IRT reading, for each test. Heart rate analysis was carried out using Kubios HRV (ver. 2.2, Biomedical Signal Analysis and Medical Imaging Group, Department of Applied Physics, University of Eastern Finland, Kuopio, Finland.). Kubios settings were adjusted in line with previous equine studies (Ille et al., 2014). Specifically, artefact correction was set to custom level 0.3, thus removing RR levels varying by more than 30% from the previous interval. This means that if a single RR interval was more than 30% different from the preceding interval, it is deemed to be an incorrect reading. Trend components were adjusted using the concept of smoothness priors set at 500ms, to avoid the effect of outlying intervals. The STD RR value, being the standard deviation

of RR intervals, was used as the HRV figure to reflect both short-term and long-term variation with the series of RR intervals. Heart rate readings for both tests were recorded for 26 subjects, allowing a comparison between handlers.

2.5 Statistical Analysis

Statistical analysis was carried out using R (R Development Core Team, 2017). Data normality was tested using Shapiro-Wilks, which indicated that data was not normally distributed. Therefore, non-parametric tests were used throughout. Wilcoxon Signed-Rank tests were used to detect potential differences in crossing time, proactivity, heart rate, heart rate variability, core temperature and discrepancy between eye temperature between familiar and unfamiliar handlers.

2.6 Ethics

Owners provided informed consent for each subject via the completion of a participant information form. All data provided was held in accordance with the Data Protection Act (1998). Both researchers and owners had the right to withdraw a subject at any time, for any reason, until the point of data analysis. Prior to commencement, the current study was authorised by the Hartpury College Ethics Committee. The authors read and abided by this journals policy on animal ethics.

3. RESULTS

There was no statistically significant difference in behaviour or any indicator of stress, depending on whether horses were handled by a familiar or unfamiliar person (Table 1).

Table 1. There were no significant differences in behaviour or physiological indicators of stress between familiar (F) and unfamiliar (UF) handlers.

Variable	n =	Handler	Median	IQR	v =	p =
Crossing Time (secs)	46	F	20.04	4.41 - 61.57	415	0.354
		UF	63.82	5.19 - 146.8		
Proactivity (%)	12	F	24.1	4.52 - 47.73	58	0.151
		UF	17.17	7.05 - 33.26		
Pre-test IRT °C	46	F	33.13	32.46 - 33.69	412	0.236
		UF	33.33	32.54 - 34.09		
Post- test IRT °C	46	F	33.15	32.54 - 33.49	440	0.388
		UF	33.08	32.3 - 33.69		
Pre-test Discrepancy °C	46	F	0.1	-0.3 - 0.7	454	0.832
		UF	0.218	-0.4 - 0.6		
Post-test Discrepancy °C	46	F	0.268	-0.2 - 0.5	411	0.373
		UF	0.1	-0.4 - 0.3		
Heart Rate	26	F	63.98	51.67 - 83.1	126	0.333
		UF	64.22	55.85 - 81.55		
Heart Rate Variability	26	F	98.79	70.71 - 143.3	163	1
		UF	98.92	80.31 - 122.9		

4. DISCUSSION

The aim of the current study was to ascertain whether a safe base effect of bonding could be observed in horses during mildly stressful handling procedures. Forty-six horses completed two novel handling tests with a familiar and unfamiliar handler. Time taken to complete the tests, proactive behaviour and physiological indicators of stress were measured. Results of the current experiment do not support the existence of a “safe base” effect of bonding in human-horse interactions (Cassidy, 1999).

Stress responses of subjects did not differ depending on whether they were handled by their owner or the unfamiliar handler. There was no difference in core eye temperature, the discrepancy in temperature between eyes, heart rate or heart rate variability. Owners care for, and train, their horses daily and, as such, are the most likely sources of human attachment. During the unfamiliar handler procedure, horses were separated from their owners and

211 presented with a potential threat, without a “safe base”. However, this does not appear to cause
212 stress in horses, indicating that neither safe base (Cassidy, 1999) nor separation distress
213 (Mikulincer and Shaver, 2003) features of bond were salient here. Time taken to complete the
214 handling tests also did not differ dependent on whether the horse was handled by their owner or
215 an unfamiliar experimental handler. In addition, there was no difference in potentially dangerous
216 proactive behaviour shown by subjects between the two handlers. This indicates horses do not
217 respond differently under situations where bonding is not possible and are not distressed at
218 being separated from their owners, even during challenging scenarios. This has implication for
219 industries such as veterinary medicine, clinical behavioural counselling and horse racing where
220 humans influence the behaviour of horses they have not interacted with previously.

221 Horses are prey animals that utilise flight to improve adaptive fitness and show consistent fear
222 responses (Lansade et al., 2008). Significant risk in horse sports and management is
223 acknowledged due to the combination of a large flight animal being routinely subjected to
224 potentially stressful procedures (Thompson et al., 2015). Some anecdotally based training
225 practices, which are often described as either “natural” or “sympathetic” horsemanship, claim
226 that bonding has benefits for resolving issues that result from these factors (Roberts, 1997).
227 They attribute reduced flight responses and improved compliance as the result of “trust”, or
228 “respect” for a leadership figure. The current experiment contradicts this and instead supports
229 previous research undermining the legitimacy of such claims (Hawson et al., 2010; McLean and
230 McGreevy, 2010). For example, it has been shown that horses will follow an unknown person
231 after “join-up” with a different individual (Krueger, 2007), or will even follow an inanimate object
232 (Henshall et al., 2012), within a round pen. In addition, the changes to behaviour resulting from
233 techniques such as round-pen interactions do not persist outside of this specialised context
234 (Krueger, 2007). Taken together, these results do not conclusively reject the possibility of bonds

between horses and their owners. They do suggest that certain features seen in fully developed attachments may not be meaningfully applied to human-horse interactions.

In the current study, the length of the relationship, the hours spent together each day and whether positive or negative reinforcement was primarily used during training was not quantified or controlled for. The type of reinforcement is known to affect subsequent reactions to humans (Sankey et al., 2010) and may therefore have confounded the current study. In addition, it is assumed that bonds take time to develop and the length of the relationship between horses and owners was not controlled for here, though it was longer than previous studies assessing the effects of familiarity (Marsbøll and Christensen, 2015). The current findings contradict those of Marsbøll and Christensen (2015), as their study noted positive effects of familiarity on handling tests. However, the subjects of that study were unusual in having only positive interaction with the familiar handler in a shorted time period. It is unlikely, despite even the best intentions, that owners in real-world scenarios can avoid any negative interactions with their horses. Despite this, a safe-base effect has been observed in human-dog relationships in which neither the length of the relationship nor the predominant training method was controlled for (INSERT MIKLOSI). This suggests that the differences between horses and dogs cannot fully be accounted for by these limitations. One key difference between Miklosi et al (2015) and the current experiment, is that subjects in the former were compared with and without any handler. In the current study, all horses were handled by the same stranger and the particular attributes of this individual are likely to affect how horses responded.

5. CONCLUSIONS

In the current study, the presence of a subject's owner did not affect behavioural or physiological indicators of stress in horses during handling tests. Results indicate that, in general, horses can be handled just as effectively without prior experience of the handler. These

findings suggest that competent handling is more salient than bond in influencing horse behaviour during handling. This has implications for industries such as veterinary practice, behaviour consultations and racing, where humans must quickly and effectively modify the behaviour of horses under potentially stressful circumstances. This experiment suggests that, in general, the presence of the horse's owner did not confer a safe-base effect. This does not conclusively reject the concept of bonds between horses and owners however. First, such bonds may be influenced by the amount and type of interaction between the dyad. It is possible that the sample tested here had not successfully developed bonds. Second, it is also possible that other features of attachment are present in human-horse interactions but that a safe base effect is not one of them. Future research into this subject is needed to explore these possibilities.

6. ACKNOWLEDGEMENTS

We sincerely thank our volunteers for their time and the use of their horses. In addition, Lauren Hunt was invaluable in sourcing and co-ordinating volunteers and Hartpury College for the use of their facilities. The authors of this manuscript have no conflict of interest to declare and no funding bodies to acknowledge.

7. AUTHORSHIP STATEMENT

The idea for this paper was conceived by Carrie Ijichi; the study was designed by Carrie Ijichi; the study was performed by Carrie Ijichi, Kym Griffin, Keith Squibb and Rebecca Favier; the data was analysed by Carrie Ijichi; the paper was written by Carrie Ijichi and edited by Kym Griffin, Keith Squibb and Rebecca Favier.

8. REFERENCES

- Cassidy, J., 1999. The nature of child's ties, in: Cassidy, J., Shaver, P. (Eds.), *Handbook of Attachment: Theory, Research and Clinical Applications*. Guilford Press, New York, pp. 3–20.
- Church, J.S., Hegadoren, P.R., Paetkau, M.J., Miller, C.C., Regev-Shoshani, G., Schaefer, A.L., Schwartzkopf-Genswein, K.S., 2014. Influence of environmental factors on infrared eye temperature measurements in cattle. *Res. Vet. Sci.* 96, 220–226. doi:10.1016/j.rvsc.2013.11.006
- De Boyer Des Roches, A., Richard-Yris, M.A., Henry, S., Ezzaouia, M., Hausberger, M., 2008. Laterality and emotions: Visual laterality in the domestic horse (*Equus caballus*) differs with objects' emotional value. *Physiol. Behav.* 94, 487–490. doi:10.1016/j.physbeh.2008.03.002
- Fallis, A., 2013. Tools of the trade or part of the family? Horses in competitive equestrian sport. *J. Chem. Inf. Model.* 53, 1689–1699. doi:10.1017/CBO9781107415324.004
- Hausberger, M., Roche, H., Henry, S., Visser, E.K., 2008. A review of the human–horse relationship. *Appl. Anim. Behav. Sci.* 109, 1–24.
- Hawson, L., McLean, A., McGreevy, P., 2010. The roles of equine ethology and applied learning theory in horse-related human injuries. *J. Vet. Behav. Clin. Appl. Res.* 5, 324–338.
- Henshall, C., Padalino, B., McGreevy, P., 2012. The radio-controlled car as herd leader? A preliminary study of escape and avoidance learning in the round-pen, in: Randle, H., Waran, N., Williams, J. (Eds.), *Proceedings of the 8th International Equitation Science Conference*. Edinburgh.
- Ijichi, C., Collins, L.M., Creighton, E., Elwood, R.W., 2013. Harnessing the power of personality assessment: Subjective assessment predicts behaviour in horses. *Behav. Processes* 96, 47–52. doi:10.1016/j.beproc.2013.02.017

308 Ille, N., Erber, R., Aurich, C., Aurich, J., 2014. Comparison of heart rate and heart rate variability
309 obtained by heart rate monitors and simultaneously recorded electrocardiogram signals in
310 nonexercising horses. *J. Vet. Behav. Clin. Appl. Res.* 9, 341–346.
311 doi:10.1016/j.jveb.2014.07.006

312 Krueger, K., 2007. Behaviour of horses in the “round pen technique.” *Appl. Anim. Behav. Sci.*
313 104, 162–170. doi:10.1016/j.applanim.2006.04.021

314 Lansade, L., Bouissou, M.-F., Erhard, H.W., 2008. Fearfulness in horses: A temperament trait
315 stable across time and situations. *Appl. Anim. Behav. Sci.* 115, 182–200.

316 Lush, J., Ijichi, C., 2018. A preliminary investigation into personality and pain in dogs. *J. Vet.*
317 *Behav.* 24, 62–68. doi:10.1016/j.jveb.2018.01.005

318 Marsbøll, A.F., Christensen, J.W., 2015. Effects of handling on fear reactions in young Icelandic
319 horses. *Equine Vet. J.* 47, 615–619. doi:10.1111/evj.12338

320 McGreevy, P., McLean, A., 2007. Roles of learning theory and ethology in equitation. *J. Vet.*
321 *Behav. Clin. Appl. Res.* 2, 108–118.

322 McLean, A.N., McGreevy, P.D.P., 2010. Horse-training techniques that may defy the principles
323 of learning theory and compromise welfare. *J. Vet. Behav. Clin. Appl. Res.* 5, 187–195.

324 Mikulincer, M., Shaver, P.R., 2003. The Attachment Behavioral System In Adulthood: Activation,
325 Psychodynamics, And Interpersonal Processes. *Adv. Exp. Soc. Psychol.* 35, 53–152.
326 doi:10.1016/S0065-2601(03)01002-5

327 Newberry, R.C., Swanson, J.C., 2008. Implications of breaking mother–young social bonds.
328 *Early Weaning* 110, 3–23.

329 Nicol, C.J., 2002. Equine learning: progress and suggestions for future research. *Equine Behav.*
330 78, 193–208.

331 Parelli, P., 1993. Natural Horsemanship. Western Horseman, Fort Worth, Texas.

332 Payne, E., Boot, M., Starling, M., Henshall, C., McLean, A., Bennett, P., McGreevy, P., 2015.
 333 Evidence of horsemanship and dogmanship and their application in veterinary contexts.
 334 Vet. J. 204, 247–254. doi:10.1016/j.tvjl.2015.04.004

335 Payne, E., DeAraugo, J., Bennett, P., McGreevy, P., 2016. Exploring the existence and potential
 336 underpinnings of dog-human and horse-human attachment bonds. Behav. Processes 125,
 337 114–121. doi:10.1016/j.beproc.2015.10.004

338 Proops, L., McComb, K., 2012. Cross-modal individual recognition in domestic horses (*Equus*
 339 *caballus*) extends to familiar humans. Proc. R. Soc. B Biol. Sci. 279, 3131–3138.
 340 doi:10.1098/rspb.2012.0626

341 R Development Core Team, 2017. R: A language and environment for statistical computing.

342 Riemer, S., Assis, L., Pike, T.W., Mills, D.S., 2016. Dynamic changes in ear temperature in
 343 relation to separation distress in dogs. Physiol. Behav. 167, 86–91.
 344 doi:10.1016/j.physbeh.2016.09.002

345 Roberts, M., 1997. The man who listens to horses. Arrow Books, London.

346 Sankey, C., Richard-Yris, M.-A., Leroy, H., Henry, S., Hausberger, M., 2010. Positive
 347 interactions lead to lasting positive memories in horses, *Equus caballus*. Anim. Behav. 79,
 348 869–875.

349 Squibb, K., Griffin, K., Favier, R., Ijichi, C., 2018. Poker Face: Discrepancies in behaviour and
 350 affective states in horses during stressful handling procedures. Appl. Anim. Behav. Sci. 0.
 351 doi:10.1016/j.applanim.2018.02.003

352 Thompson, K., McGreevy, P., McManus, P., 2015. A critical review of horse-related risk: A
 353 research agenda for safer mounts, riders and equestrian cultures. Animals 5, 561–575.

354 Travain, T., Colombo, E.S., Heinzl, E., Bellucci, D., Prato Previde, E., Valsecchi, P., 2015. Hot
355 dogs: Thermography in the assessment of stress in dogs (*Canis familiaris*)-A pilot study. *J.*
356 *Vet. Behav. Clin. Appl. Res.* 10, 17–23. doi:10.1016/j.jveb.2014.11.003

357 Walsh, F., 2009. Human-animal bonds I: The relational significance of companion animals.
358 *Fam. Process* 48, 462–480. doi:10.1111/j.1545-5300.2009.01296.x

359 Wolff, A., Hausberger, M., Le Scolan, N., 1997. Experimental tests to assess emotionality in
360 horses. *Behav. Processes* 40, 209–221.

361 Yarnell, K., Hall, C., Billett, E., 2013. An assessment of the aversive nature of an animal
362 management procedure (clipping) using behavioral and physiological measures. *Physiol.*
363 *Behav.* 118, 32–39. doi:10.1016/j.physbeh.2013.05.013

364